

Table-1: CL-HA Patients Treated & Pain Syndromes

<i>Pt. #</i>	<i>Sex</i>	<i>Age (yr.)</i>	<i>Pain Syndrome</i>	<i>Pain Duration</i>	<i>Relief Duration</i>
1	M	27	Paroxysmal LBP & Bil. buttock pain-spasm. Sciatic N.	8 yrs.	24 mos.
2	M	78	DM BLE Polyneuropathy foot pain. Tibial N.	10 yrs.	30+ mos.
2	M	82	PHN Shingles pain, Lt. ant. thigh. Ant. Femoral Cut. N.	6 mos.	3+ mos.
3	M	48	Idio. BLE Polyneuropathy foot pain. Tibial N.	11 yrs.	12 mos.
3	M	48	BLE - TTS sharp foot pain. Tibial N.	5 yrs.	12 mos.
4	F	25	Lt. Posterolat. chest wall pain, s/p crush injury. 7th ICN.	3 yrs.	12 mos.
5	F	41	Post-surg. neuritis, Lt. 5th toe. Sural Cut. N.	4 mos.	8 mos.
6	F	73	DM BLE Polyneuropathy foot pain. Tibial N.	6 mos.	2.5+ mos.
7	M	55	Lt. neck pain, C3-4, C4-5, C5-6 HNP/Cord-Root Imping.	12 mos.	6.5 mos.
7	M	55	Lt. shoulder pain, SST/Bcps td. inj. P. Supraclavicular N.	12 mos.	5.5 mos.
8	M	66	Lt. forefoot pain, DJD, digits #2, 3, MTH. Tibial N.	20 yrs.	12 mos.
9	F	44	Rt. foot post-traumatic neuritis. Sup. Peroneal N.	3.5 yrs.	4 mos.
9	F	44	Rt. LBP, SIJ osteoarthritis+osteophyte. Sup. Cluneal N.	7 yrs., 10 mos.	4 mos.
10	F	60	Idio. BLE Polyneuropathy foot pain. Tibial N.	10 yrs.	24 mos.
10	F	60	Lt. wrist pain, scaphoid osteonecrosis. Sup. Radial N.	6 mos.	3.5 mos.
10	F	60	Rt. T6-7, T7-8 pain; Intraforaminal perineural cysts. ICN	2 yrs., 5 mos.	12 mos.
11	M	45	Lt. elbow pain, post-traumatic, MVA Ulnar N.	2 yrs., 8 mos.	2.5 mos.
11	M	45	Lt. wrist pain, post-traumatic, MVA Ulnar N.	2 yrs., 8 mos.	2.5 mos.
12	F	45	Fibromyalgia, T spine. ICN-Dorsal Rami, Sensory Cut. N.	10 yrs.	4 mos.
13	M	27	Post-surg. mid-face pain, Infraorbital N.	5 yrs., 10 mos.	4 mos.
14	M	35	BLE-TTS sharp foot pain. Tibial N.	3 yrs., 10 mos.	4.5 mos.
15	F	42	Rt. Lateral Epicondylitis-Tennis Elbow. Lat. Anteb. Cut. N.	12 mos.	14 + mos.